

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>7</sup> :</b> <b>C07K 14/435, C12N 1/00, 1/19, 1/21, 5/10, 15/12, 15/63</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 00/34324</b> <b>(43) International Publication Date:</b> 15 June 2000 (15.06.00)
<b>(21) International Application Number:</b> PCT/US99/29412 <b>(22) International Filing Date:</b> 10 December 1999 (10.12.99)  <b>(30) Priority Data:</b> 09/210,330 11 December 1998 (11.12.98) US 09/444,341 19 November 1999 (19.11.99) US  <b>(71) Applicant:</b> CLONTECH LABORATORIES, INC. [US/US]; 1020 East Meadow Drive, Palo Alto, CA 94303 (US).  <b>(72) Inventors:</b> LUKYANOY, Sergey Anatolievich; ul. Golubinskaya, 13/1-161, Moscow, 117465 (RU). FRADKOV, Arcady Fedorovich; ul. Dnepropetrovskaya, 35/2-14, Moscow, 113570 (RU). LABAS, Yulii Aleksandrovich; ul. Generala Tyuleneva, 35-416, Moscow, 117465 (RU). MATZ, Mikhail Vladimirovich; ul. Teplii stan, 7/2-28, Moscow, 117465 (RU).  <b>(74) Agent:</b> ADLER, Benjamin, A.; McGregor & Adler, 8011 Candle Ln., Houston, TX 77071 (US).		<b>(81) Designated States:</b> JP, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).  <b>Published</b> <i>With international search report.</i>
<b>(54) Title:</b> FLUORESCENT PROTEINS FROM NON-BIOLUMINESCENT SPECIES OF CLASS ANTHOZOA, GENES ENCODING SUCH PROTEINS AND USES THEREOF  <b>(57) Abstract</b>  The present invention is directed to novel fluorescent proteins from non-bioluminescent organisms from the Class Anthozoa. Also disclosed are cDNAs encoding the fluorescent proteins.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

**FLUORESCENT PROTEINS FROM NON-BIOLUMINESCENT SPECIES  
OF CLASS ANTHOZOA, GENES ENCODING SUCH PROTEINS AND  
USES THEREOF**

5

**BACKGROUND OF THE INVENTION**

10

Cross-reference to Related Application

This is a divisional application of U.S.S.N. 09/210,330 filed on December 11, 1998.

15 Field of the Invention

This invention relates to the field of molecular biology. More specifically, this invention relates to novel fluorescent proteins, cDNAs encoding the proteins and uses thereof.

20 Description of the Related Art

Fluorescence labeling is a particularly useful tool for marking a protein, cell, or organism of interest. Traditionally, a protein of interest is purified, then covalently conjugated to a fluorophore derivative. For *in vivo* studies, the protein-dye complex is  
25 then inserted into cells of interest using micropipetting or a method of reversible permeabilization. The dye attachment and insertion steps, however, make the process laborious and difficult to control. An alternative method of labeling proteins of interest is to concatenate or fuse the gene expressing the protein of interest to a

gene expressing a marker, then express the fusion product. Typical markers for this method of protein labeling include  $\beta$ -galactosidase, firefly luciferase and bacterial luciferase. These markers, however, require exogenous substrates or cofactors and are therefore of limited use for *in vivo* studies.

A marker that does not require an exogenous cofactor or substrate is the green fluorescent protein (GFP) of the jellyfish *Aequorea victoria*, a protein with an excitation maximum at 395 nm, a second excitation peak at 475 nm and an emission maximum at 510 nm. GFP is a 238-amino acid protein, with amino acids 65-67 involved in the formation of the chromophore.

Uses of GFP for the study of gene expression and protein localization are discussed in detail by Chalfie et al. in *Science* 263 (1994), 802-805, and Heim et al. in *Proc. Nat. Acad. Sci.* 91 (1994), 12501-12504. Additionally, Rizzuto et al. in *Curr. Biology* 5 (1995), 635-642, discuss the use of wild-type GFP as a tool for visualizing subcellular organelles in cells, while Kaether and Gerdes in *Febs Letters* 369 (1995), 267-271, report the visualization of protein transport along the secretory pathway using wild-type GFP. The expression of GFP in plant cells is discussed by Hu and Cheng in *Febs Letters* 369 (1995), 331-334, while GFP expression in *Drosophila* embryos is described by Davis et al. in *Dev. Biology* 170 (1995), 726-729.

Crystallographic structures of wild-type GFP and the mutant GFP S65T reveal that the GFP tertiary structure resembles a barrel (Ormö et al., *Science* 273 (1996), 1392-1395; Yang, et al., *Nature Biotechnol.* 14 (1996), 1246-1251). The barrel consists of beta sheets in a compact structure, where, in the center, an alpha helix containing the chromophore is shielded by the barrel. The

compact structure makes GFP very stable under diverse and/or harsh conditions such as protease treatment, making GFP an extremely useful reporter in general. However, the stability of GFP makes it sub-optimal for determining short-term or repetitive events.

5           A great deal of research is being performed to improve the properties of GFP and to produce GFP reagents useful and optimized for a variety of research purposes. New versions of GFP have been developed, such as a "humanized" GFP DNA, the protein product of which has increased synthesis in mammalian cells (Haas, et al.,  
10 *Current Biology* 6 (1996), 315-324; Yang, et al., *Nucleic Acids Research* 24 (1996), 4592-4593). One such humanized protein is "enhanced green fluorescent protein" (EGFP). Other mutations to GFP have resulted in blue-, cyan- and yellow-green light emitting versions. Despite the great utility of GFP, however, other fluorescent proteins  
15 with properties similar to or different from GFP would be useful in the art. Novel fluorescent proteins result in possible new colors, or produce pH-dependent fluorescence. Other benefits of novel fluorescent proteins include fluorescence resonance energy transfer (FRET) possibilities based on new spectra and better suitability for  
20 larger excitation.

The prior art is deficient in novel fluorescent proteins wherein the DNA coding sequences are known. The present invention fulfills this long-standing need in the art.

25

## SUMMARY OF THE INVENTION

The present invention is directed to DNA sequences encoding fluorescent proteins selected from the group consisting of:

(a) an isolated DNA from an organism from the Class Anthozoa which encodes a fluorescent protein; (b) an isolated DNA which hybridizes to the isolated DNA of (a) and which encodes a fluorescent protein; and (c) an isolated DNA differing from the isolated DNAs of (a) and (b) in codon sequence due to the degeneracy of the genetic code and that encodes a fluorescent protein. Preferably, the DNA is isolated from a non-bioluminescent organism from Class Anthozoa. More preferably, the DNA has the sequence shown in SEQ ID No. 55 and the fluorescent protein has the amino acid sequence shown in SEQ ID No. 56.

In another embodiment of the present invention, there is provided a vector capable of expressing the DNA of the present invention in a recombinant cell comprising said DNA and regulatory elements necessary for expression of the DNA in the cell. Preferably, the DNA encodes a fluorescent protein having the amino acid sequence shown in SEQ ID No. 56.

In still another embodiment of the present invention, there is provided a host cell transfected with a vector of the present invention, such that the host cell expresses a fluorescent protein. Preferably, the cell is selected from the group consisting of bacterial cells, mammalian cells, plant cells, insect cells and yeast cells. A representative example of bacterial cell is an *E. coli* cell.

The present invention is also directed to an isolated and purified fluorescent protein coded for by DNA selected from the group consisting of: (a) isolated DNA from an organism from Class Anthozoa which encodes a fluorescent protein; (b) isolated DNA which hybridizes to the isolated DNA of (a) and which encodes a fluorescent protein; and (c) isolated DNA differing from the isolated DNAs of (a) and (b) in codon sequence due to the degeneracy of the

genetic code, and which encodes a fluorescent protein. Preferably, the protein has the amino acid sequence shown in SEQ ID No. 56.

The present invention is also directed to a DNA sequence encoding a fluorescent protein selected from the group consisting of:

5 (a) an isolated DNA which encodes a fluorescent protein, wherein said DNA is from an organism from Class Anthozoa and wherein said organism does not exhibit bioluminescence; (b) an isolated DNA which hybridizes to isolated DNA of (a) and which encodes a fluorescent protein; and (c) an isolated DNA differing from the

10 isolated DNAs of (a) and (b) in codon sequence due to degeneracy of the genetic code and which encodes a fluorescent protein. Preferably, the organism is from Sub-class Zoantharia, Order Corallimorpharia. More preferably, the organism is from Family Discosomatidae, Genus Discosoma. Even more preferably, the organism is *Discosoma sp.*

15 "*magenta*". Most particularly, the present invention is drawn to a novel fluorescent protein from *Discosoma sp.* "*magenta*", dmFP592.

The present invention is further directed to an amino acid sequence which can be used as a basis for designing an oligonucleotide probe for identification of a DNA encoding a

20 fluorescent protein by means of hybridization, wherein the amino acid sequence is selected from the group consisting of SEQ ID Nos. 3, 5, 8, 11, 12, 14. Preferably, such an oligonucleotide has a nucleotide sequence selected from the group consisting of SEQ ID Nos. 4, 6, 7, 9, 10, 13, 15, 16.

25 Other and further aspects, features, and advantages of the present invention will be apparent from the following description of the presently preferred embodiments of the invention given for the purpose of disclosure.

## BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 shows the modified strategy of 3'-RACE used to isolate the target fragments. Sequences of the oligonucleotides used are shown in Table 2. Dp1 and Dp2 are the degenerate primers used in the first and second PCR, respectively (see Tables 3 and 4 for the sequences of degenerate primers). In the case of *Discosoma sp. "magenta"*, the first degenerate primer used was NGH (SEQ ID No. 4), and the second degenerate primer used was NFP (SEQ ID No. 13). {Please confirm whether the degenerate primers used for dgFP512 are right. If not, please provide the right primers}

Figure 2 shows the excitation and emission spectrum of the novel fluorescent protein from *Discosoma sp. "magenta"*, dmFP592.

## DETAILED DESCRIPTION OF THE INVENTION

As used herein, the term "GFP" refers to the basic green fluorescent protein from *Aequorea victoria*, including prior art versions of GFP engineered to provide greater fluorescence or fluoresce in different colors. The sequence of *Aequorea victoria* GFP (SEQ ID No. 54) has been disclosed in Prasher et al., *Gene* 111 (1992), 229-33.

As used herein, the term "EGFP" refers to mutant variant of GFP having two amino acid substitutions: F64L and S65T (Heim et al., *Nature* 373 (1995), 663-664). The term "humanized" refers to changes made to the GFP nucleic acid sequence to optimize the codons



for expression of the protein in human cells (Yang et al., *Nucleic Acids Research* 24 (1996), 4592-4593).

As used herein, the term "NFP" refers to novel fluorescent protein. Specifically, "NFP" refers to dmFP592 in the present  
5 invention.

In accordance with the present invention there may be employed conventional molecular biology, microbiology, and recombinant DNA techniques within the skill of the art. Such techniques are explained fully in the literature. See, e.g., Maniatis,  
10 Fritsch & Sambrook, "Molecular Cloning: A Laboratory Manual (1982); "DNA Cloning: A Practical Approach," Volumes I and II (D.N. Glover ed. 1985); "Oligonucleotide Synthesis" (M.J. Gait ed. 1984); "Nucleic Acid Hybridization" (B.D. Hames & S.J. Higgins eds. (1985)); "Transcription and Translation" (B.D. Hames & S.J. Higgins eds.  
15 (1984)); "Animal Cell Culture" (R.I. Freshney, ed. (1986)); "Immobilized Cells and Enzymes" (IRL Press, (1986)); B. Perbal, "A Practical Guide To Molecular Cloning" (1984).

A "vector" is a replicon, such as plasmid, phage or cosmid, to which another DNA segment may be attached so as to bring about  
20 the replication of the attached segment.

A "DNA molecule" refers to the polymeric form of deoxyribonucleotides (adenine, guanine, thymine, or cytosine) in either single stranded form or a double-stranded helix. This term refers only to the primary and secondary structure of the molecule,  
25 and does not limit it to any particular tertiary forms. Thus, this term includes double-stranded DNA found, inter alia, in linear DNA molecules (e.g., restriction fragments), viruses, plasmids, and chromosomes.

A DNA "coding sequence" is a DNA sequence which is transcribed and translated into a polypeptide in vivo when placed under the control of appropriate regulatory sequences. The boundaries of the coding sequence are determined by a start codon at the 5' (amino) terminus and a translation stop codon at the 3' (carboxyl) terminus. A coding sequence can include, but is not limited to, prokaryotic sequences, cDNA from eukaryotic mRNA, genomic DNA sequences from eukaryotic (e.g., mammalian) DNA, and synthetic DNA sequences. A polyadenylation signal and transcription termination sequence may be located 3' to the coding sequence.

As used herein, the term "hybridization" refers to the process of association of two nucleic acid strands to form an antiparallel duplex stabilized by means of hydrogen bonding between residues of the opposite nucleic acid strands.

The term "oligonucleotide" refers to a short (under 100 bases in length) nucleic acid molecule.

"DNA regulatory sequences", as used herein, are transcriptional and translational control sequences, such as promoters, enhancers, polyadenylation signals, terminators, and the like, that provide for and/or regulate expression of a coding sequence in a host cell.

A "promoter sequence" is a DNA regulatory region capable of binding RNA polymerase in a cell and initiating transcription of a downstream (3' direction) coding sequence. For purposes of defining the present invention, the promoter sequence is bounded at its 3' terminus by the transcription initiation site and extends upstream (5' direction) to include the minimum number of bases or elements necessary to initiate transcription at levels detectable above background. Within the promoter sequence will be found a

transcription initiation site, as well as protein binding domains responsible for the binding of RNA polymerase. Eukaryotic promoters will often, but not always, contain "TATA" boxes and "CAT" boxes. Various promoters, including inducible promoters, may be used to  
5 drive the various vectors of the present invention.

As used herein, the terms "restriction endonucleases" and "restriction enzymes" refer to bacterial enzymes, each of which cut double-stranded DNA at or near a specific nucleotide sequence.

A cell has been "transformed" or "transfected" by  
10 exogenous or heterologous DNA when such DNA has been introduced inside the cell. The transforming DNA may or may not be integrated (covalently linked) into the genome of the cell. In prokaryotes, yeast, and mammalian cells for example, the transforming DNA may be maintained on an episomal element such as a plasmid. With respect  
15 to eukaryotic cells, a stably transformed cell is one in which the transforming DNA has become integrated into a chromosome so that it is inherited by daughter cells through chromosome replication. This stability is demonstrated by the ability of the eukaryotic cell to establish cell lines or clones comprised of a population of daughter  
20 cells containing the transforming DNA. A "clone" is a population of cells derived from a single cell or common ancestor by mitosis. A "cell line" is a clone of a primary cell that is capable of stable growth *in vitro* for many generations.

A "heterologous" region of the DNA construct is an  
25 identifiable segment of DNA within a larger DNA molecule that is not found in association with the larger molecule in nature. Thus, when the heterologous region encodes a mammalian gene, the gene will usually be flanked by DNA that does not flank the mammalian genomic DNA in the genome of the source organism. In another

example, heterologous DNA includes coding sequence in a construct where portions of genes from two different sources have been brought together so as to produce a fusion protein product. Allelic variations or naturally-occurring mutational events do not give rise to  
5 a heterologous region of DNA as defined herein.

As used herein, the term "reporter gene" refers to a coding sequence attached to heterologous promoter or enhancer elements and whose product may be assayed easily and quantifiably when the construct is introduced into tissues or cells.

10 The amino acids described herein are preferred to be in the "L" isomeric form. The amino acid sequences are given in one-letter code (A: alanine; C: cysteine; D: aspartic acid; E: glutamic acid; F: phenylalanine; G: glycine; H: histidine; I: isoleucine; K: lysine; L: leucine; M: methionine; N: asparagine; P: proline; Q: glutamine; R:  
15 arginine; S: serine; T: threonine; V: valine; W: tryptophane; Y: tyrosine; X: any residue).  $\text{NH}_2$  refers to the free amino group present at the amino terminus of a polypeptide.  $\text{COOH}$  refers to the free carboxy group present at the carboxy terminus of a polypeptide. In keeping with standard polypeptide nomenclature, *J Biol. Chem.*, 243 (1969),  
20 3552-59 is used.

The present invention is directed to an isolated DNA selected from the group consisting of: (a) isolated DNA from an organism from the Class Anthozoa which encodes a fluorescent protein; (b) isolated DNA which hybridizes to isolated DNA of (a) and  
25 which encodes a fluorescent protein; and (c) isolated DNA differing from the isolated DNAs of (a) and (b) in codon sequence due to the degeneracy of the genetic code, and which encodes a fluorescent protein. Preferably, the DNA has the sequence shown in SEQ ID No. 55

and the fluorescent protein has the amino acid sequence shown in SEQ ID No. 56.

In another embodiment of the present invention, there is provided a vector capable of expressing the DNA of the present invention in a recombinant cell comprising said DNA and regulatory elements necessary for expression of the DNA in the cell. Specifically, the DNA encodes a fluorescent protein having the amino acid sequence shown in SEQ ID No. 56.

In still another embodiment of the present invention, there is provided a host cell transfected with the vector of the present invention, which expresses a fluorescent protein of the present invention. Preferably, the cell is selected from the group consisting of bacterial cells, mammalian cells, plant cells, insect cells and yeast cells. A representative example of bacterial cell is an *E. coli* cell.

The present invention is also directed to a DNA sequence encoding a fluorescent protein selected from the group consisting of: (a) an isolated DNA which encodes a fluorescent protein, wherein said DNA is from an organism from Class Anthozoa and wherein said organism does not exhibit bioluminescence; (b) an isolated DNA which hybridizes to isolated DNA of (a) and which encodes a fluorescent protein; and (c) an isolated DNA differing from the isolated DNAs of (a) and (b) in codon sequence due to degeneracy of the genetic code and which encodes a fluorescent protein. Preferably, the organism is from Sub-class Zoantharia, Order Corallimorpharia. More preferably, the organism is from Family Discosomatidae, Genus *Discosoma*. Most preferably, the organism is *Discosoma sp. "magenta"*.

The present invention is also directed to an isolated and purified fluorescent protein coded for by DNA selected from the

group consisting of: (a) an isolated protein encoded by a DNA which encodes a fluorescent protein wherein said DNA is from an organism from Class Anthozoa and wherein said organism does not exhibit bioluminescence; (b) an isolated protein encoded by a DNA which  
5 hybridizes to isolated DNA of (a); and (c) an isolated protein encoded by a DNA differing from the isolated DNAs of (a) and (b) in codon sequence due to degeneracy of the genetic code. Preferably, the isolated and purified fluorescent protein is dmFP592.

The present invention is further directed to an amino acid  
10 sequence which can be used as a basis for designing an oligonucleotide probe for identification of a DNA encoding a fluorescent protein by means of hybridization, wherein the amino acid sequence is selected from the group consisting of SEQ ID Nos. 3, 5, 8, 11, 12, 14. Preferably, such an oligonucleotide has a nucleotide  
15 sequence selected from the group consisting of SEQ ID Nos. 4, 6, 7, 9, 10, 13, 15, 16 and is used as a primer in polymerase chain reaction. Alternatively, it can be used as a probe for hybridization screening of the cloned genomic or cDNA library.

The following examples are given for the purpose of  
20 illustrating various embodiments of the invention and are not meant to limit the present invention in any fashion.

**EXAMPLE 1****Biological Material**

Novel fluorescent proteins were identified from several  
5 genera of Anthozoa which do not exhibit any bioluminescence but  
have fluorescent color as observed under usual white light or  
ultraviolet light. Six species were chosen (see Table 1).

**TABLE 1**Anthozoa Species Used in This Study

Species	Area of Origination	Fluorescent Color
Anemonia majano	Western Pacific	bright green tentacle tips
Clavularia sp.	Western Pacific	bright green tentacles and oral disk
Zoanthus sp.	Western Pacific	green-yellow tentacles and oral disk
Discosoma sp. "red"	Western Pacific	orange-red spots oral disk
Discosoma striata	Western Pacific	blue-green stripes on oral disk
Discosoma sp. "magenta"	Western Pacific	faintly purple oral disk
Discosoma sp. "green"	Western Pacific	green spots on oral disk
Anemonia sulcata	Mediterranean	purple tentacle tips



**EXAMPLE 2****cDNA Preparation**

Total RNA was isolated from the species of interest  
5 according to the protocol of Chomczynski and Sacchi (Chomczynski  
P., et al., *Anal. Biochem.* 162 (1987), 156-159). First-strand cDNA was  
synthetized starting with 1-3 µg of total RNA using SMART PCR cDNA  
synthesis kit (CLONTECH) according to the provided protocol with the  
only alteration being that the "cDNA synthesis primer" provided in the  
10 kit was replaced by the primer TN3 (5'- CGCAGTCGACCG(T)<sub>13</sub>, SEQ ID  
No. 1) (Table 2). Amplified cDNA samples were then prepared as  
described in the protocol provided except the two primers used for  
PCR were the TS primer (5'-AAGCAGTGGTATCAACGCAGAGT, SEQ ID  
No. 2) (Table 2) and the TN3 primer (Table 2), both in 0.1 µM  
15 concentration. Twenty to twenty-five PCR cycles were performed to  
amplify a cDNA sample. The amplified cDNA was diluted 20-fold in  
water and 1 µl of this dilution was used in subsequent procedures.

**TABLE 2**Oligos Used in cDNA Synthesis and RACE

5 TN3: 5'-CGCAGTCGACCG(T)<sub>13</sub>  
(SEQ ID No. 1)

T7-TN3: 5'-GTAATACGACTCACTATAGGGCCGACGTCGACCG(T)<sub>13</sub>  
(SEQ ID No. 17)

10

TS-primer: 5'-AAGCAGTGGTATCAACGCAGAGT  
(SEQ ID No. 2)

T7-TS:  
15 5'-GTAATACGACTCACTATAGGGCAAGCAGTGGTATCAACGCAGAGT  
(SEQ ID No. 18)

T7: 5'-GTAATACGACTCACTATAGGGC  
(SEQ ID No. 19)

20

TS-oligo 5'-AAGCAGTGGTATCAACGCAGAGTACGCrGrGrG  
(SEQ ID No. 53)

25

**EXAMPLE 3****Oligo Design**

To isolate fragments of novel fluorescent protein cDNAs,  
5 PCR using degenerate primers was performed. Degenerate primers  
were designed to match the sequence of the mRNAs in regions that  
were predicted to be the most invariant in the family of fluorescent  
proteins. Four such stretches were chosen (Table 3) and variants of  
degenerate primers were designed. All such primers were directed to  
10 the 3'-end of mRNA. All oligos were gel-purified before use. Table 2  
shows the oligos used in cDNA synthesis and RACE.

**TABLE 3**

Key Amino Acid Stretches and Corresponding Degenerate Primers  
Used for Isolation of Fluorescent Proteins

5

Stretch according to A. victoria GFP (7)	Position	Amino Acid Sequence of the Key Stretch	Degenerated Primer Name and Sequence
20-25		GXVNGH (SEQ ID No. 3)	NGH: 5'- GA(C,T) GGC TGC GT(A,T,G,C) AA(T,C) GG(A,T,G) CA (SEQ ID No. 4)
31-35		GEGEG (SEQ ID No. 5)  GEGNG (SEQ ID No. 8)	GEGa: 5'- GTT ACA GGT GA(A,G) GG(A,C) GA(A,G) GG (SEQ ID No. 6) GEGb: 5'- GTT ACA GGT GA(A,G) GG(T,G) GA(A,G) GG (SEQ ID No. 7) GNGa: 5'- GTT ACA GGT GA(A,G) GG(A,C) AA(C,T) GG (SEQ ID No. 9) GNGb: 5'- GTT ACA GGT GA(A,G) GG(T,G) AA(C,T) GG (SEQ ID No. 10)
127-131		GMNFP (SEQ ID No. 11) GVNFP (SEQ ID No. 12)	NFP: 5' TTC CA(C,T) GGT (G,A)TG AA(C,T) TT(C,T) CC (SEQ ID NO. 13)
134-137		GPVM (SEQ ID No. 14)	PVMa: 5' CCT GCC (G,A)A(C,T) GGT CC(A,T,G,C) GT(A,C) ATG (SEQ ID NO. 15) PVMb: 5' CCT GCC (G,A)A(C,T) GGT CC(A,T,G,C) GT(G,T) ATG (SEQ ID NO. 16)

**EXAMPLE 4**Isolation of 3'-cDNA Fragments of nEPs

The modified strategy of 3'-RACE was used to isolate the target fragments (see Figure 1). The RACE strategy involved two consecutive PCR steps. The first PCR step involved a first degenerate primer (Table 4) and the T7-TN3 primer (SEQ ID No. 17) which has a 3' portion identical to the TN3 primer used for cDNA synthesis (for sequence of T7-TN3, Table 2). The reason for substituting the longer T7-TN3 primer in this PCR step was that background amplification which occurred when using the shorter TN3 primer was suppressed effectively, particularly when the T7-TN3 primer was used at a low concentration (0.1  $\mu$ M) (Frohman et al., (1998) *PNAS USA*, 85, 8998-9002). The second PCR step involved the TN3 primer (SEQ ID No. 1, Table 2) and a second degenerate primer (Table 4).

**TABLE 4**

Combinations of Degenerate Primers for First and Second PCR  
Resulting in Specific Amplification of 3'-Fragments of nFP cDNA

Species	First Degenerate Primer	Second Degenerate Primer
Anemonia majano	NGH (SEQ ID No. 4)	GNGb (SEQ ID No. 10)
Clavularia sp.	NGH (SEQ ID No. 4)	GEGa (SEQ ID No. 6)
Zoanthus sp.	NGH (SEQ ID No. 4)	GEGa (SEQ ID No. 6)
Discosoma sp. "red"	NGH (SEQ ID No. 4)	GEGa (SEQ ID No. 6), NFP (SEQ ID No. 13) or PVMb (SEQ ID No. 16)
Discosoma striata	NGH (SEQ ID No. 4)	NFP (SEQ ID No. 13)
Anemonia sulcata	NGH (SEQ ID No. 4)	GEGa (SEQ ID No. 6) or NFP (SEQ ID No. 13)

5

The first PCR reaction was performed as follows: 1 µl of 20-fold  
10 dilution of the amplified cDNA sample was added into the reaction  
mixture containing 1X Advantage KlenTaq Polymerase Mix with  
provided buffer (CLONTECH), 200 µM dNTPs, 0.3 µM of first

degenerate primer (Table 4) and 0.1  $\mu$ M of T7-TN3 (SEQ ID No. 17) primer in a total volume of 20  $\mu$ l. The cycling profile was (Hybaid OmniGene Thermocycler, tube control mode): 1 cycle for 95°C, 10 sec.; 55°C, 1 min.; 72°C, 40 sec; 24 cycles for 95°C, 10 sec.; 62°C, 30 sec.; 72°C, 40 sec. The reaction was then diluted 20-fold in water and 1  $\mu$ l of this dilution was added to a second PCR reaction, which contained 1X Advantage KlenTaq Polymerase Mix with the buffer provided by the manufacturer (CLONTECH), 200  $\mu$ M dNTPs, 0.3  $\mu$ M of the second degenerate primer (Table 4) and 0.1  $\mu$ M of TN3 primer.

10 The cycling profile was (Hybaid OmniGene Thermocycler, tube control mode): 1 cycle for 95°C, 10 sec.; 55°C (for GEG/GNG or PVM) or 52°C (for NFP), 1 min.; 72°C, 40 sec; 13 cycles for 95°C, 10sec.; 62°C (for GEG/GNG or PVM) or 58°C (for NFP), 30 sec.; 72°C, 40 sec. The product of PCR was cloned into PCR-Script vector (Stratagene)

15 according to the manufacturer's protocol.

Different combinations of degenerate primers were tried in the first and second PCR reactions on the DNA from each species until a combination of primers was found that resulted in specific amplification--meaning that a pronounced band of expected size

20 (about 650-800 bp for NGH and GEG/GNG and 350-500 bp for NFP and PVM--sometimes accompanied by a few minor bands) was detected on agarose gel after two PCR reactions. The primer combinations of choice for different species of the Class Anthozoa are listed in Table 4. Some other primer combinations also resulted in amplification of

25 fragments of correct size, but the sequence of these fragments showed no homology to the other fluorescent proteins identified or to *Aequorea victoria* GFP.

**EXAMPLE 5**Obtaining Full-Length cDNA Copies

Upon sequencing the obtained 3'-fragments of novel fluorescent protein cDNAs, two nested 5'-directed primers were synthesized for cDNA (Table 5), and the 5' ends of the cDNAs were then amplified using two consecutive PCRs. In the next PCR reaction, the novel approach of "step-out PCR" was used to suppress background amplification. The step-out reaction mixture contained 1x Advantage KlenTaq Polymerase Mix using buffer provided by the manufacturer (CLONTECH), 200  $\mu$ M dNTPs, 0.2  $\mu$ M of the first gene-specific primer (see Table 5), 0.02  $\mu$ M of the T7-TS primer (SEQ ID No. 18), 0.1  $\mu$ M of T7 primer (SEQ ID No. 19) and 1  $\mu$ l of the 20-fold dilution of the amplified cDNA sample in a total volume of 20  $\mu$ l. The cycling profile was (Hybaid OmniGene Thermocycler, tube control mode): 23-27 cycles for 95°C, 10 sec.; 60°C, 30 sec.; 72°C, 40 sec. The product of amplification was diluted 50-fold in water and one  $\mu$ l of this dilution was added to the second (nested) PCR. The reaction contained 1X Advantage KlenTaq Polymerase Mix with provided buffer (CLONTECH), 200  $\mu$ M dNTPs, 0.2  $\mu$ M of the second gene-specific primer and 0.1  $\mu$ M of TS primer (SEQ ID No. 2) in a total volume of 20  $\mu$ l. The cycling profile was (Hybaid OmniGene Thermocycler, tube control mode): 12 cycles for 95°C, 10 sec.; 60°C, 30 sec.; 72°C, 40 sec. The product of amplification was then cloned into pAtlas vector (CLONTECH) according to the manufacturer's protocol.



**TABLE 5**Gene-Specific Primers Used for 5'-RACE

Species	First Primer	Second (Nested) Primer
Anemonia majano	5'-GAAATAGTCAGGCATACTGGT (SEQ ID No. 20)	5'-GTCAGGCATAC TGGTAGGAT (SEQ ID No. 21)
Clavularia sp.	5'-CTTGAAATAGTCTGCTATATC (SEQ ID No. 22)	5'-TCTGCTATATC GTCTGGGT (SEQ ID No. 23)
Zoanthus sp.	5'- GTTCTTGAAATAGTCTACTATGT (SEQ ID No. 24)	5'-GTCTACTATGTCTT GAGGAT (SEQ ID No. 25)
Discosoma sp. "red"	5'-CAAGCAAATGGCAAAGGTC (SEQ ID No. 26)	5'-CGGTATTGTGGCC TTCGTA (SEQ ID No. 27)
Discosoma striata	5'-TTGTCTTCTTCTGCACAAC (SEQ ID No. 28)	5'-CTGCACAACGG GTCCAT (SEQ ID No. 29)
Anemonia sulcata	5'-CCTCTATCTTCATTCCTGC (SEQ ID No. 30)	5'-TATCTTCATTCCT GCGTAC (SEQ ID No. 31)
Discosoma sp. "magenta"	5'-TTCAGCACCCCATCACGAG (SEQ ID No. 32)	5'-ACGCTCAGAGCTG GGTTCC (SEQ ID No. 33)
Discosoma sp. "green"	5'-CCCTCAGCAATCCATCACGTTC (SEQ ID No. 34)	5'-ATTATCTCAGTGGA TGGTTC (SEQ ID No. 35)

### EXAMPLE 6

#### Expression of nFP in *E. coli*

5           To prepare a DNA construct for novel fluorescent protein expression, two primers were synthesized for each cDNA: a 5'-directed "downstream" primer with the annealing site located in the 3'-UTR of the cDNA and a 3'-directed "upstream" primer corresponding to the site of translation start site (not including the first ATG codon) (Table 6). Primers with SEQ ID Nos. 49 and 50 were the primers used to prepare the dmFP592 DNA. Both primers had 5'-heels coding for a site for a restriction endonuclease; in addition, the upstream primer was designed so as to allow the cloning of the PCR product into the pQE30 vector (Qiagen) in such a way that resulted in the fusion of reading frames of the vector-encoded 6xHis-tag and nFP. The PCR was performed as follows: 1 µl of the 20-fold dilution of the amplified cDNA sample was added to a mixture containing 1x Advantage KlenTaq Polymerase Mix with buffer provided by the manufacturer (CLONTECH), 200 µM dNTPs, 0.2 µM of upstream primer, and 0.2 µM of downstream primer, in a final total volume of 20 µl. The cycling profile was (Hybaid OmniGene Thermocycler, tube control mode): 23-27 cycles for 95°C, 10 sec.; 60°C, 30 sec.; 72°C, 40 sec. The product of this amplification step was purified by phenol-chlorophorm extraction and ethanol precipitation and then cloned into pQE30 vector using restriction endonucleases corresponding to the primers' sequence according to standard protocols.

25           All plasmids were amplified in XL-1 blue *E. coli* and purified by plasmid DNA miniprep kits (CLONTECH). The recombinant clones were selected by colony color, and grown in 3 ml of LB medium

(supplemented with 100 µg/ml of ampicillin) at 37°C overnight. 100 µl of the overnight culture was transferred into 200 ml of fresh LB medium containing 100 µg/ml of ampicillin and grown at 37°C, 200 rpm up to OD<sub>600</sub> 0.6-0.7. 1 mM IPTG was then added to the culture  
5 and incubation was allowed to proceed at 37°C for another 16 hours. The cells were harvested and recombinant protein, which incorporated 6x His tags on the N-terminus, was purified using TALON™ metal-affinity resin according to the manufacturer's protocol (CLONTECH).

10

**TABLE 6**

Primers Used to Obtain Full Coding Region of nFPs for Cloning into Expression Construct

Species	Upstream Primer	Downstream Primer
Anemonia majano	5' -acatggatccgctcttttcaaaca agtttacc (SEQ ID No. 36) BamHI	5'-tagtactcagagcttattcgta tttcagtgaatc (SEQ ID No. 37) XhoI
Clavularia sp.	L: 5'-acatggatccaacattttttga gaaacg (SEQ ID No. 38) BamHI S: 5'-acatggatccaaagctctaacc accatg (SEQ ID No. 39) BamHI	5'-tagtactcagagcaacacaa accctcagacaa (SEQ ID No. 40) XhoI
Zoanthus sp.	5'- acatggatccgctcagtc aaag cacggt (SEQ ID No. 41) BamHI	5'-tagtactcagaggttgaactacat tcttatca (SEQ ID No. 42) XhoI
Discosoma sp. "red"	5'- acatggatccaggtcttccaagaat gttacc (SEQ ID No. 43) BamHI	5'-tagtactcagaggagccaagttc agcctta (SEQ ID No. 44) XhoI
Discosoma striata	5'- acatggatccagttggtccaagagtgtg (SEQ ID No. 45) BamHI	5'-tagcagagctctatcatgcctc gtcacct (SEQ ID No. 46) SacI
Anemonia sulcata	5'-acatggatccgcttcctttttaagaagact (SEQ ID No. 47) BamHI	5'-tagtactcagagtccttgggagc ggcttg (SEQ ID No. 48) XhoI
Discosoma sp. "magenta"	5'-acatggatccagttgttccaagaatgtgat (SEQ ID No. 49) BamHI	5'-tagtactcagaggccattacg ctaacc (SEQ ID No. 50) XhoI
Discosoma sp. "green"	5'-acatggatccagtgcaacttaagaagaagaatg (SEQ ID No. 51)	5'-tagtactcagagattcggtttaat gccttg (SEQ ID No. 52)

**EXAMPLE 7**Novel Fluorescent Proteins and cDNAs Encoding the Proteins

One of the full-length cDNAs encoding fluorescent proteins  
 5 found is described herein (dmFP592). The nucleic acid sequence and  
 deduced amino acid sequence are SEQ ID Nos. 55 and 56, respectively.  
 The spectral properties of dmFP592 is listed in Table 7, and the  
 emission and excitation spectra for the dmFP592 is shown in Figure 2.

10 **TABLE 7**

Spectral Properties of the Isolated dmFP592

	Species:	Discosoma sp "magenta"	Max. Extinction Coefficient:	21,800
15	nFP Name:	dmFP592	Quantum Yield	0.09
	Absorbance Max. (nm):	573	Relative Brightness:*	0.09
20	Emission Max. (nm):	593		

\*relative brightness is extinction coefficient multiplied by quantum  
 yield divided by the same value for *A. victoria* GFP.

25 Any patents or publications mentioned in this specification  
 are indicative of the levels of those skilled in the art to which the  
 invention pertains. These patents and publications are incorporated  
 by reference to the same extent as if each individual publication was  
 specifically and individually indicated to be incorporated by  
 30 reference.

One skilled in the art will appreciate readily that the present invention is adapted to carry out the objects and obtain the ends and advantages mentioned, as well as those objects and ends inherent therein. The present examples, along with the methods, 5 procedures, treatments, molecules, and specific compounds described herein, are presently representative of preferred embodiments, are exemplary, and are not intended as limitations on the scope of the invention. Changes to the methods and compounds, and other uses, will occur to those skilled in the art and are encompassed within the 10 spirit of the invention as defined by the scope of the claims.

**WHAT IS CLAIMED IS:**

1. A DNA sequence encoding a fluorescent protein selected from the group consisting of:
  - 5 (a) an isolated DNA which encodes a fluorescent protein, wherein said DNA is from an organism from a Class Anthozoa and wherein said organism does not exhibit bioluminescence;
  - (b) an isolated DNA which hybridizes to isolated DNA of (a) above and which encodes a fluorescent protein; and
  - 10 (c) an isolated DNA differing from the isolated DNAs of (a) and (b) above in codon sequence due to degeneracy of the genetic code and which encodes a fluorescent protein.
2. The DNA sequence of claim 1, wherein said organism  
15 is from Sub-class Zoantharia.
3. The DNA sequence of claim 2, wherein said organism is from Order Corallimorpharia.
- 20 4. The DNA sequence of claim 3, wherein said organism is from Family Discosomatidae.
5. The DNA sequence of claim 4, wherein said organism is from Genus Discosoma.  
25
6. The DNA sequence of claim 5, wherein said organism is *Discosoma* sp. "magenta".

7. A DNA sequence encoding a fluorescent protein selected from the group consisting of:

(a) an isolated DNA which encodes a fluorescent protein having a nucleotide sequence shown in SEQ ID No. 55;

5 (b) an isolated DNA which hybridizes to isolated DNA of (a) above and which encodes a fluorescent protein; and

(c) an isolated DNA differing from the isolated DNAs of (a) and (b) above in codon sequence due to degeneracy of the genetic code, and which encodes a fluorescent protein.

10

8. The DNA of claim 7, wherein said DNA encodes a fluorescent protein having an amino acid sequence shown in SEQ ID No. 56.

15

9. A vector capable of expressing the DNA of claim 1 in a recombinant cell, said vector comprising said DNA of claim 1 and regulatory elements necessary for expression of the DNA in the cell.

20 10. The vector of claim 9, wherein said DNA encodes a fluorescent protein having the amino acid sequence shown in SEQ ID No. 56.

25

11. A host cell transfected with the vector of claim 9, wherein said cell is capable of expressing a fluorescent protein.

12. The host cell of claim 11, wherein said cell is selected from the group consisting of bacterial cells, mammalian cells, plant cell, yeast and insect cells.



13. The host cell of claim 12, wherein said bacterial cell is an *E. coli* cell.

14. An isolated and purified fluorescent protein coded  
5 for by DNA selected from the group consisting of:

(a) an isolated DNA which encodes a fluorescent protein from an organism from Class Anthozoa, wherein said organism does not exhibit bioluminescence;

(b) an isolated DNA which hybridizes to isolated DNA of  
10 (a) above and which encodes a fluorescent protein; and

(c) an isolated DNA differing from the isolated DNAs of (a) and (b) above in codon sequence due to degeneracy of the genetic code and which encodes a fluorescent protein.

15 15. The isolated and purified fluorescent protein of claim 14, wherein said organism is from Sub-class Zoantharia.

16. The isolated and purified fluorescent protein of  
20 claim 15, wherein said organism is from Order Corallimorpharia.

17. The isolated and purified fluorescent protein of claim 16, wherein said organism is from Family Discosomatidae.

25 18. The isolated and purified fluorescent protein of claim 17, wherein said organism is from Genus Discosoma.

19. The isolated and purified fluorescent protein of claim 18, wherein said organism is *Discosoma* sp. "*magenta*".

20. An isolated and purified fluorescent protein coded for by DNA selected from the group consisting of:

- 5 (a) isolated DNA which encodes a fluorescent protein having an amino acid sequence shown in SEQ ID No. 56;
- (b) isolated DNA which hybridizes to isolated DNA of (a) above and which encodes a fluorescent protein; and
- (c) isolated DNA differing from said isolated DNAs of (a) and (b) above in codon sequence due to degeneracy of the genetic  
10 code and which encodes a fluorescent protein.

21. The isolated and purified fluorescent protein of claim 20, wherein said protein is dmFP592.

- 15 22. An amino acid sequence which can be used as a basis for designing an oligonucleotide probe for identification of a DNA encoding a fluorescent protein by means of hybridization, wherein said sequence is selected from the group consisting of SEQ ID Nos. 3, 5, 8, 11, 12, 14.

20

23. The amino acid sequence of claim 22, wherein said oligonucleotide has a nucleotide sequence selected from the group consisting of SEQ ID Nos. 4, 6, 7, 9, 10, 13, 15, 16




FIG. 1




FIG. 2

## SEQUENCE LISTING

<110> Lukyanov, Sergey A.  
Labas, Yulii A.  
Matz, Mikhail V.  
Fradkov, Arcady F.

<120> Fluorescent Proteins from Non-Bioluminescent  
Species of Class Anthozoa, Genes Encoding Such  
Proteins and Uses Thereof

<130> D6196D9PCT

<140> 09/444,341

<141> 1999-11-19

<150> 09/210,330

<151> 1998-12-11

<160> 56

<210> 1

<211> 25

<212> DNA

<213> artificial sequence

<220>

<221> primer\_bind

<223> primer TN3 used in cDNA synthesis and RACE

<400> 1

cgcagtcgac cgtttttttt ttttt 25

<210> 2

<211> 23

<212> DNA

<213> artificial sequence

<220>

<221> primer\_bind

<223> primer TS used in cDNA synthesis and RACE

<400> 2

aagcagtggc atcaacgcag agt 23

<210> 3

<211> 6  
<212> PRT  
<213> *Aequorea victoria*  
<220>  
<222> 21  
<223> amino acid sequence of a key stretch on which  
primer NGH is based; Xaa at position 21 represents  
unknown  
<400> 3

Gly Xaa Val Asn Gly His  
5

<210> 4  
<211> 20  
<212> DNA  
<213> artificial sequence  
<220>  
<221> primer\_bind  
<222> 12  
<223> primer NGH used for isolation of fluorescent  
protein; n at position 12 represents any of the  
four bases  
<400> 4

gayggctgcg tnaayggdca

20

<210> 5  
<211> 5  
<212> PRT  
<213> *Aequorea victoria*  
<220>  
<222> 31...35  
<223> amino acid sequence of a key stretch on which  
primers GEGa and GEGb are based  
<400> 5

Gly Glu Gly Glu Gly  
5

<210> 6  
<211> 20  
<212> DNA  
<213> artificial sequence  
<220>  
<221> primer\_bind  
<223> primer GEGa used for isolation of fluorescent  
protein  
<400> 6

gttacaggtg arggmgargg 20

<210> 7  
<211> 20  
<212> DNA  
<213> artificial sequence  
<220>  
<221> primer\_bind  
<223> primer GEGb used for isolation of fluorescent  
protein  
<400> 7

gttacaggtg arggkgargg 20

<210> 8  
<211> 5  
<212> PRT  
<213> *Aequorea victoria*  
<220>  
<222> 31...35  
<223> amino acid sequence of a key stretch on which  
primers GNGa and GNGb are based  
<400> 8

Gly Glu Gly Asn Gly

5

<210> 9  
<211> 20  
<212> DNA

<213> artificial sequence  
 <220>  
 <221> primer\_bind  
 <223> primer GNGa used for isolation of fluorescent  
 protein  
 <400> 9

gttacaggtg arggmaaygg

20

<210> 10  
 <211> 20  
 <212> DNA  
 <213> artificial sequence  
 <220>  
 <221> primer\_bind  
 <223> primer GNGb used for isolation of fluorescent  
 protein  
 <400> 10

gttacaggtg arggkaaygg

20

<210> 11  
 <211> 5  
 <212> PRT  
 <213> *Aequorea victoria*  
 <220>  
 <222> 127..131  
 <223> amino acid sequence of a key stretch on which  
 primer NFP is based  
 <400> 11

Gly Met Asn Phe Pro

5

<210> 12  
 <211> 5  
 <212> PRT  
 <213> *Aequorea victoria*  
 <220>  
 <222> 127..131



<223> amino acid sequence of a key stretch on which  
primer NFP is based

<400> 12

Gly Val Asn Phe Pro

5

<210> 13

<211> 20

<212> DNA

<213> artificial sequence

<220>

<221> primer\_bind

<223> primer NFP used for isolation of fluorescent  
protein

<400> 13

ttccayggtr tgaayttycc

20

<210> 14

<211> 4

<212> PRT

<213> *Aequorea victoria*

<220>

<222> 134...137

<223> amino acid sequence of a key stretch on which  
primers PVMa and PVMb are based

<400> 14

Gly Pro Val Met

<210> 15

<211> 21

<212> DNA

<213> artificial sequence

<220>

<221> primer\_bind

<222> 15

<223> primer PVMa used for isolation of fluorescent protein; n at position 15 represents any of the four bases

<400> 15

cctgccrayg gtcnngtmat g

21

<210> 16

<211> 21

<212> DNA

<213> artificial sequence

<220>

<221> primer\_bind

<222> 15

<223> primer PVMb used for isolation of fluorescent protein; n at position 15 represents any of the four bases

<400> 16

cctgccrayg gtcnngtkat g

21

<210> 17

<211> 47

<212> DNA

<213> artificial sequence

<220>

<221> primer\_bind

<223> primer T7-TN3 used in cDNA synthesis and RACE

<400> 17

gtaatacgac tcactatagg gccgcagtcg accgtttttt tttttt

47

<210> 18

<211> 45

<212> DNA

<213> artificial sequence

<220>

<221> primer\_bind

<223> primer T7-TS used in cDNA synthesis and RACE

<400> 18

gtaatacgac tcactatagg gcaagcagtg gtatcaacgc agagt 45

<210> 19  
<211> 22  
<212> DNA  
<213> artificial sequence  
<220>  
<221> primer\_bind  
<223> primer T7 used in cDNA synthesis and RACE  
<400> 19

gtaatacgac tcactatagg gc 22

<210> 20  
<211> 21  
<212> DNA  
<213> artificial sequence  
<220>  
<221> primer\_bind  
<223> gene-specific primer used for 5'-RACE for *Anemonia majano*  
<400> 20

gaaatagtca ggcatactgg t 21

<210> 21  
<211> 20  
<212> DNA  
<213> artificial sequence  
<220>  
<221> primer\_bind  
<223> gene-specific primer used for 5'-RACE for *Anemonia majano*  
<400> 21

gtcaggcata ctggtaggat 20

<210> 22

<211> 21  
<212> DNA  
<213> artificial sequence  
<220>  
<221> primer\_bind  
<223> gene-specific primer used for 5'-RACE for  
*Clavularia sp.*  
<400> 22

cttgaaatag tctgctatat c 21

<210> 23  
<211> 19  
<212> DNA  
<213> artificial sequence  
<220>  
<221> primer\_bind  
<223> gene-specific primer used for 5'-RACE for  
*Clavularia sp.*  
<400> 23

tctgctatat cgtctgggt 19

<210> 24  
<211> 23  
<212> DNA  
<213> artificial sequence  
<220>  
<221> primer\_bind  
<223> gene-specific primer used for 5'-RACE for *Zoanthus*  
*sp.*  
<400> 24

gttcttgaaa tagtctacta tgt 23

<210> 25  
<211> 20  
<212> DNA

<213> artificial sequence  
<220>  
<221> primer\_bind  
<223> gene-specific primer used for 5'-RACE for *Zoanthus*  
sp.  
<400> 25

gtctactatg tcttgaggat

20

<210> 26  
<211> 19  
<212> DNA  
<213> artificial sequence  
<220>  
<221> primer\_bind  
<223> gene-specific primer used for 5'-RACE for  
*Discosoma* sp. "red"  
<400> 26

caagcaaattg gcaaaggtc

19

<210> 27  
<211> 19  
<212> DNA  
<213> artificial sequence  
<220>  
<221> primer\_bind  
<223> gene-specific primer used for 5'-RACE for  
*Discosoma* sp. "red"  
<400> 27

cggtattgtg gccttcgta

19

<210> 28  
<211> 19  
<212> DNA  
<213> artificial sequence  
<220>  
<221> primer\_bind

<223> gene-specific primer used for 5'-RACE for  
*Discosoma striata*

<400> 28

ttgtcttctt ctgcacaac

19

<210> 29

<211> 17

<212> DNA

<213> artificial sequence

<220>

<221> primer\_bind

<223> gene-specific primer used for 5'-RACE for  
*Discosoma striata*

<400> 29

ctgcacaacg ggtccat

17

<210> 30

<211> 20

<212> DNA

<213> artificial sequence

<220>

<221> primer\_bind

<223> gene-specific primer used for 5'-RACE for *Anemonia*  
*sulcata*

<400> 30

cctctatctt catttcctgc

20

<210> 31

<211> 20

<212> DNA

<213> artificial sequence

<220>

<221> primer\_bind

<223> gene-specific primer used for 5'-RACE for *Anemonia*  
*sulcata*

<400> 31

tatcttcatt tcctgcgtac

20

<210> 32  
<211> 19  
<212> DNA  
<213> artificial sequence  
<220>  
<221> primer\_bind  
<223> gene-specific primer used for 5'-RACE for  
*Discosoma* sp. "magenta"  
<400> 32

ttcagcaccc catcacgag

19

<210> 33  
<211> 19  
<212> DNA  
<213> artificial sequence  
<220>  
<221> primer\_bind  
<223> gene-specific primer used for 5'-RACE for  
*Discosoma* sp. "magenta"  
<400> 33

acgctcagag ctgggttcc

19

<210> 34  
<211> 22  
<212> DNA  
<213> artificial sequence  
<220>  
<221> primer\_bind  
<223> gene-specific primer used for 5'-RACE for  
*Discosoma* sp. "green"  
<400> 34

ccctcagcaa tccatcacgt tc

22

<210> 35

<211> 20  
<212> DNA  
<213> artificial sequence  
<220>  
<221> primer\_bind  
<223> gene-specific primer used for 5'-RACE for  
*Discosoma sp. "green"*  
<400> 35

attatctcag tggatggttc 20

<210> 36  
<211> 31  
<212> DNA  
<213> artificial sequence  
<220>  
<221> primer\_bind  
<223> upstream primer used to obtain full coding region  
of nFPs from *Anemonia majano*  
<400> 36

acatggatcc gctctttcaa acaagtttat c 31

<210> 37  
<211> 34  
<212> DNA  
<213> artificial sequence  
<220>  
<221> primer\_bind  
<223> downstream primer used to obtain full coding  
region of nFPs from *Anemonia majano*  
<400> 37

tagtactcga gcttattcgt atttcagtga aatc 34

<210> 38  
<211> 29  
<212> DNA  
<213> artificial sequence  
<220>



<221> primer\_bind  
<223> upstream primer used to obtain full coding region  
of nFPs from *Clavularia sp.*  
<400> 38

acatggatcc aacatttttt tgagaaacg 29

<210> 39  
<211> 28  
<212> DNA  
<213> artificial sequence  
<220>  
<221> primer\_bind  
<223> upstream primer used to obtain full coding region  
of nFPs from *Clavularia sp.*  
<400> 39

acatggatcc aaagctctaa ccaccatg 28

<210> 40  
<211> 31  
<212> DNA  
<213> artificial sequence  
<220>  
<221> primer\_bind  
<223> downstream primer used to obtain full coding  
region of nFPs from *Clavularia sp.*  
<400> 40

tagtactcga gcaacacaaa ccctcagaca a 31

<210> 41  
<211> 28  
<212> DNA  
<213> artificial sequence  
<220>  
<221> primer\_bind  
<223> upstream primer used to obtain full coding region  
of nFPs from *Zoanthus sp.*

<400> 41  
acatggatcc gctcagtcaa agcacggt 28

<210> 42  
<211> 32  
<212> DNA  
<213> artificial sequence  
<220>  
<221> primer\_bind  
<223> downstream primer used to obtain full coding  
region of nFPs from *Zoanthus sp.*  
<400> 42

tagtactcga ggttggaact acattcttat ca 32

<210> 43  
<211> 31  
<212> DNA  
<213> artificial sequence  
<220>  
<221> primer\_bind  
<223> upstream primer used to obtain full coding region  
of nFPs from *Discosoma sp.* "red"  
<400> 43

acatggatcc aggtcttcca agaatgttat c 31

<210> 44  
<211> 29  
<212> DNA  
<213> artificial sequence  
<220>  
<221> primer\_bind  
<223> downstream primer used to obtain full coding  
region of nFPs from *Discosoma sp.* "red"  
<400> 44

tagtactcga ggagccaagt tcagcctta 29

<210> 45  
<211> 28  
<212> DNA  
<213> artificial sequence  
<220>  
<221> primer\_bind  
<223> upstream primer used to obtain full coding region  
of nFPs from *Discosoma striata*  
<400> 45

acatggatcc agttggtcca agagtgtg 28

<210> 46  
<211> 28  
<212> DNA  
<213> artificial sequence  
<220>  
<221> primer\_bind  
<223> downstream primer used to obtain full coding  
region of nFPs from *Discosoma striata*  
<400> 46

tagcgagctc tatcatgcct cgtcacct 28

<210> 47  
<211> 31  
<212> DNA  
<213> artificial sequence  
<220>  
<221> primer\_bind  
<223> upstream primer used to obtain full coding region  
of nFPs from *Anemonia sulcata*  
<400> 47

acatggatcc gcttcctttt taaagaagac t 31

<210> 48  
<211> 28  
<212> DNA  
<213> artificial sequence

&lt;220&gt;

&lt;221&gt; primer\_bind

<223> downstream primer used to obtain full coding  
region of nFPs from *Anemonia sulcata*

&lt;400&gt; 48

tagtactcga gtccttggga gcggttg

28

&lt;210&gt; 49

&lt;211&gt; 30

&lt;212&gt; DNA

&lt;213&gt; artificial sequence

&lt;220&gt;

&lt;221&gt; primer\_bind

<223> upstream primer used to obtain full coding region  
of nFPs from *Discosoma sp. "magenta"*

&lt;400&gt; 49

acatggatcc agttgttcca agaatgtgat

30

&lt;210&gt; 50

&lt;211&gt; 26

&lt;212&gt; DNA

&lt;213&gt; artificial sequence

&lt;220&gt;

&lt;221&gt; primer\_bind

<223> downstream primer used to obtain full coding  
region of nFPs from *Discosoma sp. "magenta"*

&lt;400&gt; 50

tagtactcga ggccattacg ctaatc

26

&lt;210&gt; 51

&lt;211&gt; 31

&lt;212&gt; DNA

&lt;213&gt; artificial sequence

&lt;220&gt;

&lt;221&gt; primer\_bind

<223> upstream primer used to obtain full coding region  
of nFPs from *Discosoma sp. "green"*

<400> 51  
 acatggatcc agtgactta aagaagaaat g 31

<210> 52  
 <211> 29  
 <212> DNA  
 <213> artificial sequence  
 <220>  
 <221> primer\_bind  
 <223> downstream primer used to obtain full coding  
 region of nFPs from *Discosoma* sp. "green"  
 <400> 52

tagtactcga gattcggttt aatgccttg 29

<210> 53  
 <211> 33  
 <212> DNA  
 <213> artificial sequence  
 <220>  
 <221> primer\_bind  
 <223> TS-oligo used in cDNA synthesis and RACE  
 <400> 53

aagcagtggc atcaacgcag agtacgcrgr grg 33

<210> 54  
 <211> 238  
 <212> PRT  
 <213> *Aequorea victoria*  
 <220>  
 <223> amino acid sequence of GFP  
 <400> 54

Met	Ser	Lys	Gly	Glu	Glu	Leu	Phe	Thr	Gly	Val	Val	Pro	Ile	Leu
			5						10					15
Val	Glu	Leu	Asp	Gly	Asp	Val	Asn	Gly	His	Lys	Phe	Ser	Val	Ser
			20						25					30
Gly	Glu	Gly	Glu	Gly	Asp	Ala	Thr	Tyr	Gly	Lys	Leu	Thr	Leu	Lys
			35						40					45

Phe Ile Cys Thr Thr Gly Lys Leu Pro Val Pro Trp Pro Thr Leu		
	50	55 60
Val Thr Thr Phe Ser Tyr Gly Val Gln Cys Phe Ser Arg Tyr Pro		
	65	70 75
Asp His Met Lys Gln His Asp Phe Phe Lys Ser Ala Met Pro Glu		
	80	85 90
Gly Tyr Val Gln Glu Arg Thr Ile Phe Phe Lys Asp Asp Gly Asn		
	95	100 105
Tyr Lys Thr Arg Ala Glu Val Lys Phe Glu Gly Asp Thr Leu Val		
	110	115 120
Asn Arg Ile Glu Leu Lys Gly Ile Asp Phe Lys Glu Asp Gly Asn		
	125	130 135
Ile Leu Gly His Lys Leu Glu Tyr Asn Tyr Asn Ser His Asn Val		
	140	145 150
Tyr Ile Met Ala Asp Lys Gln Lys Asn Gly Ile Lys Val Asn Phe		
	155	160 165
Lys Ile Arg His Asn Ile Glu Asp Gly Ser Val Gln Leu Ala Asp		
	170	175 180
His Tyr Gln Gln Asn Thr Pro Ile Gly Asp Gly Pro Val Leu Leu		
	185	190 195
Pro Asp Asn His Tyr Leu Ser Thr Gln Ser Ala Leu Ser Lys Asp		
	200	205 210
Pro Asn Glu Lys Arg Asp His Met Val Leu Leu Glu Phe Val Thr		
	215	220 225
Ala Ala Gly Ile Thr His Gly Met Asp Glu Leu Tyr Lys		
	230	235

<210> 55  
 <211> 876  
 <212> DNA  
 <213> *Discosoma sp. "magenta"*  
 <220>  
 <221> CDS  
 <223> cDNA sequence of dmFP592  
 <400> 55

agtttcagcc agtgacaggg tgagctgccca ggtattctaa caagatgagt 50  
 tgttccaaga atgtgatcaa ggagttcatg aggttcaagg ttcgtatgga 100

```

aggaacgggc aatgggcacg agtttgaaat aaaaggcgaa ggtgaagga 150
ggccttacga aggtcactgt tccgtaaagc ttatggtaac caagggtgga 200
cctttgccat ttgcttttga tattttgtca ccacaatttc agtatggaag 250
caaggtatat gtcaaacacc ctgccgacat accagactat aaaaagctgt 300
catttcctga gggattttaa tgggaaaggg tcatgaactt tgaagacggg 350
ggcgtgggta ctgtatccca agattccagt ttgaaagacg gctgtttcat 400
ctacgagggtc aagttcattg ggggtgaactt tccttctgat ggacctgtta 450
tgcagaggag gacacggggc tgggaagcca gctctgagcg tttgtatcct 500
cgtgatgggg tgctgaaagg agacatccat atggctctga ggctggaagg 550
aggcgggccat tacctcgttg aattcaaaag tatttacatg gtaaagaagc 600
cttcagtgcg gttgccaggc tactattatg ttgactccaa actggatatg 650
acgagccaca acgaagatta cacagtcgtt gagcagtatg aaaaaacca 700
gggacgccac catccgttca ttaagcctct gcagtgaact cggctcagtc 750
atggattagc ggtaatggcc acaaaaggca cgatgatcgt tttttaggaa 800
tgcagccaaa aattgaaggt tatgacagta gaaatacaag caacaggctt 850
tgcttattaa acatgtaatt gaaaac 876

```

```

<210>      56
<211>      235
<212>      PRT
<213>      Discosoma sp. "magenta"
<220>
<223>      amino acid sequence of dmFP592
<400>      56

```

```

Met Ser Cys Ser Lys Asn Val Ile Lys Glu Phe Met Arg Phe Lys
      5                      10                      15

Val Arg Met Glu Gly Thr Val Asn Gly His Glu Phe Glu Ile Lys
      20                      25                      30

Gly Glu Gly Glu Gly Arg Pro Tyr Glu Gly His Cys Ser Val Lys
      35                      40                      45

Leu Met Val Thr Lys Gly Gly Pro Leu Pro Phe Ala Phe Asp Ile
      50                      55                      60

Leu Ser Pro Gln Phe Gln Tyr Gly Ser Lys Val Tyr Val Lys His
      65                      70                      75

Pro Ala Asp Ile Pro Asp Tyr Lys Lys Leu Ser Phe Pro Glu Gly
      80                      85                      90

```

Phe Lys Trp Glu Arg Val Met Asn Phe Glu Asp Gly Gly Val Val		
100	105	110
Thr Val Ser Gln Asp Ser Ser Leu Lys Asp Gly Cys Phe Ile Tyr		
115	120	125
Glu Val Lys Phe Ile Gly Val Asn Phe Pro Ser Asp Gly Pro Val		
130	135	140
Met Gln Arg Arg Thr Arg Gly Trp Glu Ala Ser Ser Glu Arg Leu		
145	150	155
Tyr Pro Arg Asp Gly Val Leu Lys Gly Asp Ile His Met Ala Leu		
160	165	170
Arg Leu Glu Gly Gly Gly His Tyr Leu Val Glu Phe Lys Ser Ile		
175	180	185
Tyr Met Val Lys Lys Pro Ser Val Gln Leu Pro Gly Tyr Tyr Tyr		
190	195	200
Val Asp Ser Lys Leu Asp Met Thr Ser His Asn Glu Asp Tyr Thr		
205	210	215
Val Val Glu Gln Tyr Glu Lys Thr Gln Gly Arg His His Pro Phe		
220	225	230
Ile Lys Pro Leu Gln		
235		



## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US99/29412

## A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) :C07K 14/435; C12N 1/00, 1/19, 1/21, 5/10, 15/12, 15/63

US CL :Please See Extra Sheet.

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 435/325, 326, 348, 410, 252.3, 252.33, 254.11, 320.1; 530/350, 855; 536/23.5

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

Please See Extra Sheet.

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X, P	MATZ et al. Fluorescent proteins from non-bioluminescent Anthozoa species. Nature Biotechnology. October 1999, Vol. 17, No. 10, pages 969-973, entire document.	1-23
L	MACEK et al. Intrinsic tryptophan fluorescence of equinatoxin II, a pore forming polypeptide from the sea anemone, <i>Actinia equina</i> L, monitors its interaction with lipid membranes. European Journal of Biochemistry. 1995, Vol. 234, No. 1, pages 329-335, entire document. (cited as L document because it establishes that equinatoxin II fluoresces.)	1-2, 9, 11-13
---		-----
X		14-16

BEST AVAILABLE COPY



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
*A* document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
*E* earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
*I* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*G* document member of the same patent family
*O* document referring to an oral disclosure, use, exhibition or other means	
*P* document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

03 FEBRUARY 2000

Date of mailing of the international search report

23 FEB 2000

Name and mailing address of the ISA/US  
Commissioner of Patents and Trademarks  
Box PCT  
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

GABRIELE ELISABETH BUGAISKY

Telephone No. (703) 308-0196

## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US99/29412

## C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	LORENZ et al. Isolation and expression of a complementary DNA encoding <i>Renilla reniformis</i> luciferase. Proc. Natl. Acad. Sci., USA. May 1991, Vol. 88, No. 10, pages 4438-4442, entire document.	1-23
X	ANDERLUH et al. Cloning, sequencing, and expression of equinatoxin II. Biochem. Biophys. Res. Commun. 18 March 1996, Vol. 220, No. 2, pages 437-442, entire document.	1-2, 9, 11-15

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US99/29412

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:
  
2. ☒ Claims Nos.: 7-8, 10, 20-23  
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:  
  
The sequence diskette containing the CRF that was submitted with the application was found to be defective. Thus, no search of the specifically recited sequences could be carried out. The search report is based entirely on a word search.
  
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
  
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
  
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
  
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.  
☐ No protest accompanied the payment of additional search fees.

## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US99/29412

### A. CLASSIFICATION OF SUBJECT MATTER:

US CL :

435/325, 326, 348, 410, 252.3, 252.33, 254.11, 320.1; 530/350; 536/23.5

### B. FIELDS SEARCHED

Electronic data bases consulted (Name of data base and where practicable terms used):

WEST-files USPT, DWPI, EPAB, JPAB; Dialog files Medline, Biosis, Dissertation Abstracts, Scisearch, Biological Records, STN-CAS files registry, caplus  
search terms: Cnidar?, anthozo?, anemone?, coral, zoanth?, corallimorph? discosom?, rhodactis, magenta, protein?, fluoresc?, bioluminesc?, actinodisc?